

Module 5

We will be matching pre and post-tests to assess the integration of new knowledge, skills and attitudes with respect to working with adolescents who are at risk or actively using substances. As such, we will not ask for your name, but a unique identifier which will allow us to pair the pre with the post-test. Please fill in your unique identifier below - it should be your birthdate (Date_Month_Year), gender (M/F) and your first and last initial. For example, for a female named Jane Doe who was born on February 3, 1980, her unique identifier would be **03021980FJD**.

Unique Identifier: _____

Please circle one of the following.

PRE-TEST

POST-TEST

Answer the following true/false questions.

1. Adolescent substance use must be identified and addressed as soon as possible
2. Staying in treatment for an adequate period of time and continuity of care afterward are not important.
3. Role of the family is pivotal
4. Treatment must be general
5. Family members can play an important role in treatment engagement and in treatment outcomes. Research has shown that family interventions can bring about

Fill in the blanks on the following questions:

6. Effective _____ for adolescents requires some form of behavioral therapy.

7. Name three different types of treatment.

8. Name three areas of training needs to develop treatment professionals.

9. Name three different types of family treatment

Check the box which most closely corresponds to your response.

| | Strongly Agree | Agree | Neither Agree nor Disagree | Disagree | Strongly Disagree |
|---|-----------------------|--------------|-----------------------------------|-----------------|--------------------------|
| Effectively treating AOD use in adolescents requires also identifying and treating other mental | | | | | |

health conditions they might have.

The largest proportion of adolescents who receive treatment are referred from juvenile justice system.

The effects on families may continue for generations.

The therapist has to be willing to challenge their habitual ways of engaging family members into the process of treatment.